Study of Seasonal Variation of Biomass and Seaweed Composition in Tidal Areas of Bushehr Province (North Part of the Persian Gulf)

Dadolahi-Sohrab, Ali^{1*}; Garavand, Mohsen²; Emadabadi, Abolfazl³

- 1- Assistant Professor, Department of Marine Environment, Faculty of Marine Natural Resources, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran. E-mail: p dadolahi@yahoo.com
- 2- M.Sc. Department of Marine Biology, Faculty of Marine Science and Oceanography, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran. E-mail: mgk garavand@yahoo.com
- 3- M.Sc. Department of Marine Biology, Faculty of Marine Science and Oceanography, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran. E-mail: a_emadabady@yahoo.com

Received Date: July 04, 2010 *Corresponding Author Accepted Date: December 31, 2011

© 2012 Oceanography All rights reserved.

Abstract

The study was carried out to evaluate the seasonal variations of dominant seaweed biomass and species composition at six different sites during 4 seasons from autumn 2008 till summer 2009 in the sand flats of tidal areas of the Bushehr Province. Algal samples were collected using quadrate sampling (0.5 × 0.5 m) from 4 different tidal areas (upper mid littoral, mid mid littoral, low mid littoral and infra littoral) at each site. Samples collected were sorted as far as possible at the location and placed in plastic bags. The plastic bags were placed in an ice-chest during transportation to the laboratory. Algal samples were distinguished using valid references. In sum, this study recorded 13 dominant seaweed species including 3 Chlorophyta, 4 Phaeophyta and 6 Rhodophyta were. Among Chlorophyta, Entromorpha intestinalis was the dominant species at all sites. This is while, Cystoseira myrica and Gracilaria corticatay both showed maximum biomass among the Phaeophyta and Rhodophyta, respectively.

Keywords: Seaweed, Tidal areas, Biomass, Persian Gulf.